



Delaware Weekly Influenza Report

MMWR Week 51 (December 17 - 23, 2017)

Delaware Division of Public Health

National Influenza Synopsis 2017-2018:

National data are updated Friday of each week. Please visit <http://www.cdc.gov/flu/weekly/> for the most current information. During MMWR Week 51 (December 17 – 23, 2017) influenza activity increased sharply in the United States. The most frequently identified influenza virus subtype reported by public health laboratories during week 51 was influenza A(H3). The percentage of respiratory specimens testing positive for influenza in clinical laboratories increased. **Widespread influenza activity** was reported by 36 states (Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Texas, Virginia, Washington, West Virginia, Wisconsin and Wyoming). **Regional influenza activity** was reported by Puerto Rico and 13 states (Alaska, Hawaii, Iowa, Maine, Michigan, Nevada, New Hampshire, New Jersey, North Carolina, Rhode Island, Tennessee, Utah and Vermont). **Local influenza activity** was reported by Delaware. The District of Columbia, the U.S. Virgin Islands and Guam did not report. Both national and state data are provisional and subject to change as additional reports are received.

Delaware Influenza Surveillance 2017-2018:

During MMWR Week 51, there were 96 laboratory-confirmed cases of influenza reported among Delaware residents, bringing the total to 225 confirmed cases for the 2017-2018 season. Reports of influenza-like illness (ILI) received from participating providers, facilities and institutions in Delaware show ILI is 0.50% compared with Delaware’s 2017-2018 baseline of 2.0%. Nationally, ILI is 5.0% compared with the 2017-2018 national baseline of 2.2%.

Level of Influenza Activity in Delaware, MMWR Week 51:

Local	Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in a single region of the state.
Influenza-like illness (ILI) is defined as patients presenting with fever of 100° F or greater, cough and/or sore throat in the absence of a known cause other than influenza.	
<p>No Activity: No laboratory-confirmed cases of influenza and no reported increase in the number of cases of ILI.</p> <p>Sporadic: Small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of ILI.</p> <p>Local: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in a single region of the state.</p> <p>Regional: <i>Outbreaks of influenza or increases in ILI and recent laboratory-confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions.</i>³</p> <p>Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.</p>	

¹ 2017-2018 Region 3 (DE, DC, MD, PA, VA and WV) baseline = 2.0%.

² Laboratory-confirmed case = case confirmed by viral culture or PCR.

³ Region = population under surveillance in a defined geographical subdivision of a state. Regions typically include several counties. Regional doesn't apply to states with ≤ four counties.

<u>In this report:</u>	<u>page</u>
Table 1a. Influenza positive cases reported statewide and county, by subtype / lineage and MMWR week, Delaware 2017-18	2
Table 1b. Influenza positive cases reported statewide and county, by age group and MMWR week, Delaware 2017-18	2
Figure 1. Confirmed cases of influenza by type and subtype / lineage, by MMWR week, Delaware 2017-18	3
Table 2. Influenza-related hospitalizations, statewide and county, by age group and MMWR week, Delaware 2017-18	3
Table 3. Influenza-related deaths by MMWR week, Delaware 2017-18	3
Table 4. Numbers of influenza cases reported by flu season, Delaware 2004-05 through 2017-18	4
Figure 2. Percentage of visits for influenza-like illness reported by sentinel providers participating in CDC's ILINet, Delaware 2017-18	4
Figure 3. Influenza-like illness reported by ILI reporting partners by MMWR week, Delaware 2017-18	5
Figure 4a. Percentage of emergency department (ED) visits due to ILI/Flu by MMWR week, Delaware 2017-18	5
Figure 4b. County-specific percentages of ED visits due to ILI/Flu by MMWR week, Delaware 2017-18	5
Summary of International Influenza Activity	6

Table 1a. Influenza positive¹ cases reported² statewide and county by subtype (A) or lineage (B)³, Delaware 2017-18

Confirmed Flu Cases by Subtype / Lineage		Week 40	Week 41	Week 42	Week 43	Week 44	Week 45	Week 46	Week 47	Week 48	Week 49	Week 50	Week 51	YTD	YTD Total	YTD County %
STATEWIDE	A / 2009 H1N1	0	0	0	1	0	0	0	1	1	1	2	3	9	225	
	A / 2012 H3N2	0	0	1	0	1	0	2	1	6	15	9	9	44		
	A / no subtype	0	0	0	4	2	4	7	4	6	17	30	61	135		
	B / Yamagata	0	0	0	0	0	0	0	0	1	0	3	7	11		
	B / Victoria	0	0	0	0	0	0	0	0	0	0	0	0	0		
	B / no lineage	0	0	0	0	1	0	1	0	2	2	4	16	26		
New Castle County	A / 2009 H1N1	0	0	0	1	0	0	0	0	1	1	1	0	4	96	42.7%
	A / 2012 H3N2	0	0	1	0	0	0	2	1	5	13	9	7	38		
	A / no subtype	0	0	0	1	0	1	3	3	2	7	2	12	31		
	B / Yamagata	0	0	0	0	0	0	0	0	1	0	2	6	9		
	B / Victoria	0	0	0	0	0	0	0	0	0	0	0	0	0		
	B / no lineage	0	0	0	0	1	0	1	0	0	1	1	10	14		
Kent County	A / 2009 H1N1	0	0	0	0	0	0	0	1	0	0	0	2	3	45	20.0%
	A / 2012 H3N2	0	0	0	0	0	0	0	0	0	2	0	0	2		
	A / no subtype	0	0	0	3	1	2	2	1	2	2	8	16	37		
	B / Yamagata	0	0	0	0	0	0	0	0	0	0	0	0	0		
	B / Victoria	0	0	0	0	0	0	0	0	0	0	0	0	0		
	B / no lineage	0	0	0	0	0	0	0	0	0	0	2	1	3		
Sussex County	A / 2009 H1N1	0	0	0	0	0	0	0	0	0	0	1	1	2	84	37.3%
	A / 2012 H3N2	0	0	0	0	1	0	0	0	1	0	0	2	4		
	A / no subtype	0	0	0	0	1	1	2	0	2	8	20	33	67		
	B / Yamagata	0	0	0	0	0	0	0	0	0	0	1	1	2		
	B / Victoria	0	0	0	0	0	0	0	0	0	0	0	0	0		
	B / no lineage	0	0	0	0	0	0	0	0	2	1	1	5	9		

Table 1b. Influenza positive¹ cases reported² statewide and county by age group, Delaware 2017-18


Confirmed Flu Cases by Age Group		Week 40	Week 41	Week 42	Week 43	Week 44	Week 45	Week 46	Week 47	Week 48	Week 49	Week 50	Week 51	YTD	YTD Total	YTD County %
STATEWIDE	0-4 years	0	0	0	0	1	1	0	0	1	4	3	10	20	225	
	5-24 years	0	0	0	0	1	0	1	2	2	7	9	18	40		
	25-49 years	0	0	0	3	0	0	2	0	3	5	5	20	38		
	50-64 years	0	0	0	0	0	1	3	2	5	4	14	13	42		
	65+ years	0	0	1	2	2	2	4	2	5	15	17	35	85		
New Castle County	0-4 years	0	0	0	0	0	1	0	0	0	2	2	5	10	96	42.7%
	5-24 years	0	0	0	0	0	0	1	1	2	4	3	8	19		
	25-49 years	0	0	0	1	0	0	2	0	1	4	0	8	16		
	50-64 years	0	0	0	0	0	0	1	2	2	1	6	6	18		
	65+ years	0	0	1	1	1	0	2	1	4	11	4	8	33		
Kent County	0-4 years	0	0	0	0	0	0	0	0	0	0	0	2	2	45	20.0%
	5-24 years	0	0	0	1	1	0	0	1	0	0	1	6	9		
	25-49 years	0	0	0	1	0	0	0	0	0	1	2	2	7		
	50-64 years	0	0	0	0	0	1	1	0	2	2	3	2	11		
	65+ years	0	0	0	1	0	1	1	1	0	1	4	7	16		
Sussex County	0-4 years	0	0	0	0	1	0	0	0	1	2	1	3	8	84	37.3%
	5-24 years	0	0	0	0	0	0	0	0	0	3	5	4	12		
	25-49 years	0	0	0	0	0	0	0	0	2	0	3	10	15		
	50-64 years	0	0	0	0	0	0	1	0	1	1	5	5	13		
	65+ years	0	0	0	0	1	1	1	0	1	3	9	20	36		

¹ Based on patients with positive nucleic acid or viral culture test results reported to the Division of Public Health.

² Reports are by the date the laboratory results are obtained. As a result, prior weeks' counts may be adjusted to reflect additional cases received.

³ The Division of Public Health Laboratory now has the capability to identify lineage for Influenza B. Since some laboratories in the state do not have this capability, those influenza cases will be categorized as Influenza B, no lineage identified.

Figure 1. Confirmed cases¹ of influenza by type and subtype/lineage, Delaware 2017-18*



¹ Based on patients with positive nucleic acid or viral culture test results reported to the Delaware Division of Public Health.
Data Source: Season 2017 - 2018 Influenza Positive Specimens from Delaware, Reported by WHO/NREVSS Collaborating Laboratories

Table 2. Influenza-related hospitalizations statewide and county, by age group, Delaware 2017-18

Hospitalized Flu Cases by Age Group		Week 40	Week 41	Week 42	Week 43	Week 44	Week 45	Week 46	Week 47	Week 48	Week 49	Week 50	Week 51	YTD	YTD Total	YTD County %
STATEWIDE	0-4 years	0	0	0	0	0	0	0	0	0	0	1	1	2	63	
	5-24 years	0	0	0	0	0	0	0	0	0	1	1	0	2		
	25-49 years	0	0	0	0	0	0	0	0	0	2	0	1	3		
	50-64 years	0	0	0	0	0	0	0	1	3	1	8	2	15		
	65+ years	0	0	1	2	1	1	1	2	3	11	9	10	41		
New Castle County	0-4 years	0	0	0	0	0	0	0	0	0	0	0	1	1	38	60.3%
	5-24 years	0	0	0	0	0	0	0	0	0	1	0	0	1		
	25-49 years	0	0	0	0	0	0	0	0	0	2	0	1	3		
	50-64 years	0	0	0	0	0	0	0	1	2	0	3	2	8		
	65+ years	0	0	1	1	1	0	1	1	3	10	3	4	25		
Kent County	0-4 years	0	0	0	0	0	0	0	0	0	0	0	0	0	11	17.5%
	5-24 years	0	0	0	0	0	0	0	0	0	0	0	0	0		
	25-49 years	0	0	0	0	0	0	0	0	0	0	0	0	0		
	50-64 years	0	0	0	0	0	0	0	0	1	1	2	0	4		
	65+ years	0	0	0	1	0	1	0	1	0	0	2	2	7		
Sussex County	0-4 years	0	0	0	0	0	0	0	0	0	0	1	0	1	14	22.2%
	5-24 years	0	0	0	0	0	0	0	0	0	0	1	0	1		
	25-49 years	0	0	0	0	0	0	0	0	0	0	0	0	0		
	50-64 years	0	0	0	0	0	0	0	0	0	0	3	0	3		
	65+ years	0	0	0	0	0	0	0	0	0	1	4	4	9		

Table 3. Influenza-related deaths, Delaware 2017-18

Influenza-Related Deaths	Week 40	Week 41	Week 42	Week 43	Week 44	Week 45	Week 46	Week 47	Week 48	Week 49	Week 50	Week 51	YTD
	0	0	0	0	0	0	0	0	0	1	0	1	2


Table 4. Annual number of influenza cases reported by flu season, Delaware 2004-05 through 2017-18

Influenza Season	Total Annual Influenza Cases
2004 – 2005	995
2005 – 2006	541
2006 – 2007	508
2007 – 2008	1,401
2008 – 2009	738
2009 – 2010	2,247
2010 – 2011	1,479
2011 – 2012	267
2012 – 2013	1,781
2013 – 2014	1,842
2014 – 2015	2,390
2015 – 2016	1,842
2016 – 2017	4,590
2017 – 2018 (YTD)	225

U.S. Outpatient Influenza-Like Illness Surveillance Network (ILINet) Sentinel Providers

An ILINet (sentinel) provider conducts surveillance for influenza-like illness (ILI) in collaboration with the Division of Public Health and the Centers for Disease Control and Prevention (CDC). Data reported by ILINet providers, in combination with other influenza surveillance data, provide a national and statewide picture of influenza activity in the U.S.


Figure 2. Percentage of visits for influenza-like illness reported by sentinel providers¹ participating in the U.S. Outpatient ILI Surveillance Network (ILINet), Delaware 2017-18



¹ Twelve of 13 sentinel providers reported.

² Regional baseline is calculated by CDC using non-influenza weeks from the previous three influenza seasons. Delaware is in Region 3 that also includes DC, MD, PA, VA and WV.

Figure 3. Influenza-like illness reported by influenza surveillance ILI reporting partners¹, Delaware 2017-18



- ¹ ILINet reporting partners include long-term care facilities, colleges / universities, correctional facilities and daycare facilities.
- ² % ILI= percentage of residents with ILI symptoms. Seven long-term care facilities reported.
- ³ % ILI= percentage of student visits for ILI; One university reported.
- ⁴ % ILI= percentage of visits for ILI at the correctional facility; Nine correctional facilities reported.
- ⁵ % ILI= percentage of children absent with ILI; One daycare provider reported.

Figure 4a. Percentage of emergency department (ED) visits due to ILI/Flu by MMWR Week, Delaware 2017-18





Figure 4b. County-specific percentages of ED visits due to ILI/Flu by MMWR Week, Delaware 2017-18



Summary of International Influenza Activity

Worldwide, influenza A(H3N2) and B viruses accounted for the majority of influenza detections. Influenza activity continued to increase in the temperate zone of the northern hemisphere while in the temperate zone of the southern hemisphere activity decreased at inter-seasonal levels. In Central America and the Caribbean, influenza activity remained low.

In North America, overall influenza activity continued to increase in the region, with detections of predominantly influenza A(H3N2) viruses.

In the Caribbean and Central American countries, respiratory illness indicators and influenza activity remained low in general but respiratory syncytial virus (RSV) activity remained high in several countries.

In Europe, influenza activity continued to increase, but remained low in most of the countries, with detections of predominantly influenza B followed by influenza A(H3N2) viruses.

In Western Asia, elevated levels of influenza activity were reported in recent weeks, with influenza A(H1N1)pdm09 predominantly detected. In Central Asia, low to no influenza activity was reported. In East Asia, influenza activity remained low in most of the countries with the exception of China where influenza like illness (ILI) and influenza percentage positive continued to increase, with influenza B Yamagata-lineage viruses predominantly detected. In South East Asia, low levels of influenza activity were reported. In Southern Asia, influenza activity remained low in general. Detections of influenza A(H1N1)pdm09 and A(H3N2) viruses were reported in India and of all seasonal subtypes in the Islamic Republic of Iran.

In Northern Africa, low levels of influenza activity were reported. Detections of influenza A(H1N1)pdm09 virus increased slightly in Tunisia. In Western Africa, influenza virus detections were reported in Burkina Faso, Ghana, and Sierra Leone, with influenza A(H1N1) pdm09 virus predominating. In Middle Africa, sporadic detections of influenza A were reported in Cameroon. In Eastern Africa, influenza A(H3N2) and B detections were reported in Madagascar and Mozambique.

In the tropical countries of South America, influenza and RSV activity remained at low levels overall.

In the temperate zone of the Southern Hemisphere, influenza activity decreased overall to inter-seasonal levels.

Reference: World Health Organization (WHO), 2017. Influenza Update number 305 (12/25/17). Retrieved December 29, 2017, from http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/. Reports are updated biweekly.

NOTE: Data provided do not reflect the total number of individuals who have been infected with the influenza virus in Delaware during the reporting period due to the following factors:

- Many people ill with influenza-like symptoms do not seek medical care.
- Many who do seek medical care are not tested for influenza.
- The Delaware Public Health Laboratory is limited by capacity to processing a maximum of three specimens per day from each reporting entity.

The Delaware Division of Public Health (DPH) is committed to serving you better by providing the most accurate, up-to-date influenza data available.

- For general information on influenza, visit flu.delaware.gov or <http://dhss.delaware.gov/dhss/dph/dpc/immunize-flu.html>.
- For specific information on DPH flu clinics, visit <http://dhss.delaware.gov/dhss/dph/fluclinics.html>.
- For questions on Delaware's weekly flu report, call the DPH Office of Infectious Disease Epidemiology: 302-744-4990.
- For questions regarding influenza vaccination, please call 302-744-1060.