

FACTS *for* FAMILIES

No. 73

December 2009

Self-Injury in Adolescents

Self-injury is the act of deliberately destroying body tissue, at times to change a way of feeling. Self-injury is seen differently by groups and cultures within society. This appears to have become more popular lately, especially in adolescents. The causes and severity of self-injury can vary. Some forms may include:

- carving
- scratching
- branding
- marking
- picking, and pulling skin and hair
- burning/abrasions
- cutting
- biting
- head banging
- bruising
- hitting
- tattooing
- excessive body piercing

Some adolescents may self-mutilate to take risks, rebel, reject their parents' values, state their individuality or merely be accepted. Others, however, may injure themselves out of desperation or anger to seek attention, to show their hopelessness and worthlessness, or because they have suicidal thoughts. These children may suffer from serious psychiatric problems such as depression, psychosis, Posttraumatic Stress Disorder (PTSD) and Bipolar Disorder. Additionally, some adolescents who engage in self-injury may develop Borderline Personality Disorder as adults. Some young children may resort to self-injurious acts from time to time but often grow out of it. Children with mental retardation and/or autism as well as children who have been abused or abandoned may also show these behaviors.

Why do adolescents self-injure?

Self-injury is a complex behavior and symptom that results from a variety of factors. Adolescents who have difficulty talking about their feelings may show their emotional tension, physical discomfort, pain and low self-esteem with self-injurious behaviors. Although some teenagers may feel like the steam in the pressure cooker has been released following the act of harming themselves, others may feel hurt, anger, fear and hate. The effects of peer pressure and contagion can also influence adolescents to injure themselves. Even though fads come and go, most of the wounds on the adolescents' skin

will be permanent. Occasionally, teenagers may hide their scars, burns and bruises due to feeling embarrassed, rejected or criticized about their physical appearance.

What can parents and teenagers do about self-injury?

Parents are encouraged to talk with their children about respecting and valuing their bodies. Parents should also serve as role models for their teenagers by not engaging in acts of self-harm. Some helpful ways for adolescents to avoid hurting themselves include learning to:

- accept reality and find ways to make the present moment more tolerable.
- identify feelings and talk them out rather than acting on them.
- distract themselves from feelings of self-harm (for example, counting to ten, waiting 15 minutes, saying "NO!" or "STOP!," practicing breathing exercises, journaling, drawing, thinking about positive images, using ice and rubber bands)
- stop, think, and evaluate the pros and cons of self-injury.
- soothe themselves in a positive, non-injurious, way.
- practice positive stress management.
- develop better social skills.

Evaluation by a mental health professional may assist in identifying and treating the underlying causes of self-injury. Feelings of wanting to die or kill themselves are reasons for adolescents to seek professional care emergency. A child and adolescent psychiatrist can also diagnose and treat the serious psychiatric disorders that may accompany self-injurious behavior.

For additional information see *Facts for Families*:

[#4 Depression](#)

[#38 Manic Depressive Illness](#)

[#70 PTSD](#)

[#52 Comprehensive Psychiatric Evaluation](#)

[#66 Helping Teenagers with Stress](#)

[#5 Child Abuse](#)

[#10 Teen Suicide](#)

If you find *Facts for Families*® helpful and would like to make good mental health a reality, consider donating to the [Campaign for America's Kids](#). Your support will help us continue to produce and distribute Facts for Families, as well as other vital mental health information, free of charge.

You may also mail in your contribution. Please make checks payable to the AACAP and send to *Campaign for America's Kids*, P.O. Box 96106, Washington, DC 20090.

The American Academy of Child and Adolescent Psychiatry (AACAP) represents over 8,500 child and adolescent psychiatrists who are physicians with at least five years of additional training beyond medical school in general (adult) and child and adolescent psychiatry.

Facts for Families© information sheets are developed, owned and distributed by AACAP. Hard copies of *Facts* sheets may be reproduced for personal or educational use without written permission, but cannot be included in material presented for sale or profit. All *Facts* can be viewed and printed from the AACAP

Self-Injury, “Facts for Families,” No. 73 (12/09)

website (www.aacap.org). *Facts* sheets may not be reproduced, duplicated or posted on any other website without written consent from AACAP. Organizations are permitted to create links to AACAP’s website and specific *Facts* sheets. For all questions please contact the AACAP Communications & Marketing Coordinator, ext. 154.

If you need immediate assistance, please dial 911.

Copyright © 2012 by the American Academy of Child and Adolescent Psychiatry.